

We are grateful to A. B. Harris and J. Kukula for helpful conversations. One of us (T.C.L.) is grateful to the Alfred P. Sloan Foundation for financial assistance. This work was supported in part by National Science Foundation Grant No. DMR 76-21073, National Science Foundation, Materials Research Laboratory Grant No. DMR 76-00678, and U. S. Office of Naval Research Grant No. N000 14-76-C-106.

^(a)Permanent address: The Racah Institute of Physics, The Hebrew University, Jerusalem, Israel.

¹G. Toulouse, *Commun. Phys.* **2**, 115 (1977); J. Villain, *J. Phys. C* **10**, 1717 (1977).

²S. Kirkpatrick, *Phys. Rev. B* **16**, 4630 (1977); J. Vaniminen, *J. Phys. C* **10**, L537 (1977).

³J. E. Dzyaloshinskii and G. E. Volovik, *J. Phys. (Paris)* **39**, 693 (1978).

⁴A. Aharony, Y. Imry, and S.-k. Ma, *Phys. Rev. Lett.* **37**, 1364 (1976); A. P. Young, *J. Phys. C* **10**, L257 (1977); G. Grinstein, *Phys. Rev. Lett.* **37**, 944 (1976).

⁵R. Harris, M. Plischke, and M. J. Zuckerman, *Phys. Rev. Lett.* **31**, 160 (1973); J. H. Chen and T. C. Lubensky, *Phys. Rev. B* **16**, 2106 (1977); A. Aharony, *Phys. Rev. B* **12**, 1038 (1975); E. Pytte, to be published.

⁶R. A. Pelcovits, E. Pytte, and J. Rudnick, *Phys. Rev. Lett.* **40**, 476 (1978); A. Aharony, to be published.

⁷P. G. de Gennes, *The Physics of Liquid Crystals* (Clarendon Press, Oxford, England, 1974), Chap. 4.

⁸G. Toulouse and M. Kléman, *J. Phys. (Paris), Lett.* **37**, L149 (1976); M. Kléman, L. Michel, and G. Toulouse, *J. Phys. (Paris), Lett.* **38**, L195 (1977); N. D. Mermin, unpublished lecture notes.

⁹S. F. Edwards and P. W. Anderson, *J. Phys. F* **5**, 965 (1975).

Search for Linear Polarization of the Cosmic Background Radiation

Phil M. Lubin and George F. Smoot

Space Sciences Laboratory and Lawrence Berkeley Laboratory,

University of California, Berkeley, California 94720

(Received 6 October 1978)

We present preliminary measurements of the linear polarization of the cosmic microwave background (3 K blackbody) radiation. These ground-based measurements are made at 9-mm wavelength. We find no evidence for linear polarization, and set an upper limit for a polarized component of 0.8 mK with 95% confidence level. This implies that the present rate of expansion of the Universe is isotropic to 1 part in 10^6 , assuming no reionization of the primordial plasma after recombination.

The observed cosmic microwave background radiation is generally thought to be a relic of the hot, dense, initial phase of the Universe. This radiation should then have characteristics which reflect its thermal origin and the geometry of the early Universe. The simple hot big-bang model predicts that this radiation has a blackbody (Planckian) spectrum, is unpolarized, exhibits the statistical properties of blackbody radiation,¹ and is isotropic in a reference frame comoving with the expansion of the Universe.

The spectrum of this radiation has been extensively investigated.² The best experiment to date is that of Woody and Richards.³ Although their results qualitatively follow a blackbody spectrum, they report a 5σ deviation (σ is 1 standard deviation) from the best-fit blackbody curve. There have also been many measurements of the angular distribution of this radiation with recent reports of a (3.5 ± 0.6) -mK-amplitude first-order

(cosine) anisotropy.^{4,5} It is thought that this anisotropy is a result of the Doppler shift caused by the motion of the solar system relative to the cosmic radiation. If this interpretation is correct, the cosmic radiation shows an intrinsic anisotropy of less than 1 part in 3000 (1 mK). The polarization properties of the radiation have been largely unexplored although, in their original paper on the discovery of the background radiation, Penzias and Wilson assert that this radiation is "within the limits of our observations... unpolarized...".⁶ Nanos⁷ and Caderni *et al.*⁸ have searched for evidence of linear polarization with negative results.

It is of great interest to measure the polarization because of its potential to detect and distinguish deviations from the simple big-bang model. Rees has shown that any intrinsic anisotropy in the cosmic radiation present at the time of decoupling or last scattering manifests itself as a

linear polarization through the mechanism of Thomson scattering.⁹ Similarly, a last scattering by matter with a large velocity in the comoving frame produces a net polarization. The predicted polarization anisotropy is highly model dependent, but is estimated to be on the order of one to one-hundredth of the intrinsic intensity anisotropy.

Recent reports of the first-order anisotropy and the deviations from a blackbody spectrum provide added stimulus for investigating the polarization. We report here preliminary measurements made at 9-mm wavelength covering three circles at northern declinations δ of 38° , 53° , and 63° .

The polarimeter is shown schematically in Fig. 1. The antenna is a dual-mode, corrugated horn having nearly equal gain patterns for both linear polarizations.¹⁰ Below the antenna is a Faraday-rotation switch which alternately rotates the incoming planes of polarization from $+45^\circ$ to -45° at 100 Hz. Between the switch and the receiver is an element which selects one of the linear polarizations. Therefore, the input signal to the receiver switches rapidly between two orthogonal linear polarizations. The output of the receiver is analyzed for a signal synchronous with the switching of the Faraday rotator. Thus, the apparatus measures the difference in power between the two orthogonal polarization states. The information is digitized and recorded on magnetic

tape. In addition, the polarimeter physically rotates in 45° increments around the beam axis at 100-sec intervals. This rotation allows us to measure both orthogonal components of any possible polarized signal and provides a mean for distinguishing between external signals and imbalances in the apparatus.

The antenna (7° beam width) has very low side lobes, such that the measured response in all directions more than 90° from the antenna axis is reduced by a factor of 10^8 over the forward gain. The addition of a ground shield further suppresses the effects of Earth, so that the total intensity received from Earth is less than 0.2 mK. We estimate the apparent polarized signal resulting from this ground pickup to be less than 0.05 mK. The Faraday-rotation switch is a ferrite device and thus is sensitive to temperature and magnetic fields. The switch is temperature controlled to 0.2°C and shielded against external magnetic fields. The ferrite has a temperature-dependent absorption coefficient of about $5\text{ mK}/^\circ\text{C}$. Thermal mass insures that the ferrite temperature does not change faster than $0.3^\circ\text{C}/\text{h}$. Hence even without the small correction we make in the data analysis, the ambient temperature changes could not contribute more than a 0.08-mK signal for our 100-sec integration time. The ferrite also has a magnetization-dependent absorption coefficient. Tests indicate that the effect of Earth's field is less than 0.08 mK at the 95% confidence level. The receiver is a double-balanced mixer with a sensitivity of $100\text{ mK}/\text{Hz}^{1/2}$, a center frequency of 33.1 GHz, and a bandwidth of 500 MHz.

A polarized source calibrates the system. The source consists of a grid of copper strips on a thin plastic sheet and an ambient blackbody source. Independent laboratory measurements and theoretical calculations show that the calibrator is nearly an ideal polarizer and produces a polarized signal whose amplitude (280 K) is the difference between the ambient blackbody source (295 K) and the sky (15 K).

The measurements were made at the Lawrence Berkeley Laboratory (38° north latitude) during June, July, and August of 1978. The data at 38° declination were obtained by pointing the instrument vertically while the 53° and 63° declination data were taken by tilting north 15° and 25° , respectively.

We analyzed the data by first calculating two Stokes parameters, $Q = T_{NS} - T_{EW}$ and $U = T_{NW, SE} - T_{NE, SW}$. Thus, for example, Q is the difference




FIG. 1. Schematic of polarimeter.




FIG. 2. Measured Stokes parameters Q and U as a function of position in the sky. Data eliminated because of side-lobe contamination by the sun are indicated.

in temperature of the north-south and east-west polarizations. Q and U are signal-averaged into hourly bins with a 24-sidereal-hour period. The largest source of extraneous polarized radiation is expected to be synchronous radiation from the Galaxy. We have used data from polarization surveys at 820 MHz,^{11,12} combined with extrapolations of low-frequency intensity measurements, to estimate the polarized signal at 33 GHz. The polarized component of the Galactic synchrotron signal should be less than 0.05 ± 0.04 mK over all the areas observed. Statistical tests of the data show that the fluctuations are consistent with those expected from receiver noise.

We therefore conclude that the signal-averaged data shown in Fig. 2 are a good measure of any

possible cosmic signal. The data are fitted to a constant (dc) level and periods of 12 and 24 h, excluding the sun-contaminated portion of the vertical ($\delta = 38^\circ$) data. We find no evidence for linear polarization in the cosmic background radiation and set the 95%-confidence-level limits shown in Table I. Table II summarizes the current status of polarization measurements.

Since all measurements are consistent with no or very little polarization, one must consider the possibility that depolarization mechanisms destroy intrinsic polarization. Our calculations indicate that Faraday rotation, the birefringence of a plasma in a magnetic field, is small for the Galaxy and for the ionosphere. A universal magnetic field of magnitude greater than 10^{-7} G and an electron density of 10^{-5} cm⁻³ could effectively depolarize a signal, though there is no evidence that such fields exist. Measurements¹³ of the polarization of extragalactic objects, particularly quasars with red shifts $z \approx 2$ indicate that depolarization through Faraday rotation at 33 GHz is not important to at least a red shift $z \geq 10$. Brans¹⁴ has shown that the anisotropic expansion itself may tend to rotate the polarization, through the metric anisotropies, but this effect has been shown to be small.^{8,15}

The null result observed is consistent with a thermal origin for the cosmic background radiation and an isotropic expansion. Using the calculations of Rees and assuming recombination at $z \sim 1500$ with no subsequent reheating, our limit of 0.8 mK implies a present differential expansion limit of about 1 part in 10^6 . If the primordial plasma is reionized after recombination, the limit on the present differential expansion rate is about 1 part in 10^4 for a critically dense Universe. The present intensity anisotropy roughly equals the Hubble-expansion anisotropy at the last scat-

TABLE I. Limits on polarization anisotropies at 95% confidence level (in millikelvin). The limits are the magnitudes of the measured values plus 2 standard deviations. The sky roughness is the rms fluctuation on a 7° scale size.

Declination	38°		53°		63°	
Period	Q	U	Q	U	Q	U
dc	0.3	0.6	0.8	0.6	0.4	1.1
24 h	0.8	0.7	0.4	0.8	0.8	0.8
12 h	0.4	0.6	0.5	0.7	0.8	0.9
Sky roughness	0.6		0.8		0.7	
Days of data	30		10		8	

TABLE II. Measured limits on linear polarization at 95% confidence level.

Reference	Wavelength (cm)	Sky coverage (deg)	Limit on Q/T and U/T (%)
Penzias and Wilson (Ref. 6)	7.35	Scattered	10
Nanos (Ref. 7)	3.2	$\delta = +40$	0.05
This experiment	0.9	$\delta = +38$	0.03
		$\delta = +53$	0.03
		$\delta = +63$	0.04
Caderni <i>et al.</i> (Ref. 8)	0.05 - 0.3	Near Galactic center	0.1-1

tering, while the polarization anisotropy is the value produced during the last few scatterings.

In the case of no reionization after recombination, the 1-mK limit⁵ on the intensity anisotropy implies a more stringent limit on the present differential expansion rate whereas in the case of reheating after recombination a comparable limit is obtained. The limits on the rms roughness of the sky imply that gravitational waves with wavelengths which are large (greater than few hundred megaparsecs) on a cosmic scale have an energy density which is much lower than is needed to close the Universe.⁸

This work was supported in part by the U. S. Department of Energy and the National Aeronautics and Space Administration. One of us (P.L.) gratefully acknowledges support from the National Science Foundation as a National Science Foundation Fellow. We are indebted to C. Witebsky for calculating the Galactic backgrounds and to L. W. Alvarez, J. S. Aymong, A. Buffington, L. Deutch, L. Drexler, H. B. Dougherty, J. H. Gibson, M. V. Gorenstein, T. S. Mast, R. A. Muller, and H. J. Yamada for support and valuable suggestions.

¹R. Hanbury Brown and R. Q. Twiss, *Nature* (London) **178**, 1447 (1956).

²L. Danese and G. De Zotti, *Nuovo Cimento* **7**, 3, 277 (1977).

³D. Woody and P. Richards, to be published.

⁴B. E. Corey and D. T. Wilkinson, *Bull. Astron. Astrophys. Soc.* **8**, 351 (1976).

⁵G. F. Smoot, M. V. Gorenstein, and R. A. Muller, *Phys. Rev. Lett.* **39**, 14, 898 (1977).

⁶A. A. Penzias and R. W. Wilson, *Astrophys. J.* **142**, 419 (1965).

⁷G. P. Nanos, Ph.D. thesis, Princeton University, 1974 (unpublished).

⁸N. Caderni, R. Fabbi, B. Melchiorri, F. Melchiorri, and V. Natale, *Phys. Rev. D* **17**, 1901 (1978).

⁹M. J. Rees, *Astrophys. J.* **153**, L1 (1968).

¹⁰M. Janssen, S. Bednarczyk, S. Gulkis, H. Marlin, and G. F. Smoot, "IEEE, Antennas and Propagation" (to be published).

¹¹E. M. Berkhuijsen, *Astron. Astrophys. Suppl.* **5**, 263 (1972).

¹²E. M. Berkhuijsen, *Astron. Astrophys.* **14**, 359 (1971).

¹³F. Gardner, D. Morris, and J. Whiteoak, *Aust. J. Phys.* **22**, 79 (1969).

¹⁴C. Brans, *Astrophys. J.* **197**, 1 (1975).

¹⁵A. M. Anile and R. A. Breuer, *Astrophys. J.* **217**, 353 (1977).